GENETIC DISTANCE AND TRANSMISSION OF GENETIC INFORMATION AT RAPD AND SSR LOCI IN REGULARLY AND LATE BUDDING PEDUNCULATE OAK (QUERCUS ROBUR L.)

Oliver Werner & Gunter M. Rothe*

Institut für Allgemeine Botanik, Fachbereich 21 (Biologie), Johannes Gutenberg-Universität, Saarstrasse 21, D–55099 Mainz, Germany; *corresponding author

Received March 21, 2002; accepted December 18, 2002

ABSTRACT

One late budding and four regularly budding stands of pedunculate oak (*Quercus robur* L.) which were planted in the 19^{th} century in the south-west of Germany were investigated with respect to their genetic variabilities and genetic distances at 36 RAPD (random amplified DNA) and 28 SSR (simple sequence repeat) loci. The late budding stand and one of the regularly budding stands are licenced for seed harvesting. Offspring of these stands were included in the investigations.

Among the seven populations, subpopulation differentiation – based on the Tanimoto-distance – was largest for one of the regularly budding stands as was the expected heterozygosity. On the other hand, subpopulation differentiation and expected heterozygosity were lowest for offspring of the late budding stand.

At seven loci in offspring of the late budding stand and at four loci in offspring of the regularly budding stand, locus frequencies deviated from the corresponding parental population by more than 30 %. Subpopulation differentiation in offspring of the late budding population and the regularly budding population were respectively 3.51 % and 1.46 % larger than in the corresponding parental population. The mean expected heterozygosity in offspring of the late budding parent stand and the regularly budding stand decreased by 5.36 and 2.93 %. The different transmission of genetic information in the regularly and the late budding population is discussed in terms of different adaptational processes and seed management.

Key words: budding, heterozygosity, offspring, pedunculate oak, population genetics, *Quercus robur* L., Slovenian pedunculte oak, Tanimoto distance

INTRODUCTION

Since the Middle Ages pedunculate oak (Quercus robur L.) was often planted in Germany, regardless of its habitat demands (RÖHRIG & GUSSONE 1980). One of the reasons was that a considerable part of its natural habitat, especially river valley woodlands, had been destroyed. In the Southwest of Germany, in the state of Rhineland-Palatinate, pedunculate oak is therefore mostly found outside its original habitat. When these stands were created at the end of the 19th century they were mostly planted with seedlings of unknown origin. On the other hand, on some elevated areas of the Eifel hills, seeds of late budding varieties, imported from Slovenia, were used to establish pedunculate oak woods. These populations bud one to three weeks later than indigenous ones, which start budding at the beginning of May. The Slavic varieties were planted because they are resistent to late frosts and several parasites such as oak Tortrix species and winter moth species. Some of the regularly budding and the late budding stands are currently licensed for seed harvesting. In years of

© ARBORA PUBLISHERS

abundant fructification acorns are collected from these stands and seedlings raised for two to three years in a nursery before they are used for reforestations.

This type of forest management raises some questions about the genetics of these artificially established populations. For example: how large are their genetic variations and distances and how much genetic information is transmitted from parent to offspring populations?

We tried to answer these questions through the use of RAPD (random amplified polymorphic DNA) and SSR (simple sequence repeat) markers. RAPD markers have been applied to the study of several tree species (AIDE & RIVERA 1998, ALLNUTT *et al.* 1998, CHAL-MERS *et al.* 1994, GILLIES *et al.* 1999, YEH *et al.* 1995) including oak (PETIT *et al.* 1993). Nuclear microsatellite markers were studied by applying AMP (anchored microsatellite primed) PCR (polymerase chain reaction) technology (ZIETKIEWICZ *et al.* 1994). Both markers are dominant markers but they directly assess DNA variation without any need for DNA sequencing information (NEWTON *et al.* 1999, WINTER & KAHL 1995).

MATERIALS AND METHODS

Stand locations

Investigations were performed on pedunculate oak in the state of Rhineland-Palatinate (Rheinland-Pfalz), Southwest-Germany. Trees were planted in the 19th century and are growing in the forest districts of Hagenbach, Mayen, Simmern, Speyer and Trippstadt (Table 1, Figure 1). At each stand, 50 oaks which had been classified before according to their leaf and fruit traits as morphologically pure pedunculate oak (*Quercus robur* L.) (ROMMEL *et al.* 1995) were probed. In addition, 50 offspring, each at the age of eight years were studied, which originated from the investigated stands in the forest districts of Hagenbach and Mayen which are licenced for seed harvesting. The offspring are part of a provenance trial which had been established at forest district Neupfalz (Figure 1, Table 1).

Mode of sampling

Short twigs, with recently flushed leaves, were shot

from the trees, transported to the lab and kept in water until extracted within the next two days. From the samplings young leaves were taken and stored between wet filter papers at 8 °C and extracted within three days.

DNA extraction

Leaves were washed with distilled water and dabbed between layers of filter paper. DNA was extracted by a modified version of DELLAPORTA *et al.* (1983) and Doyle (1991): Approximately 1 gram of fresh leaves was ground under liquid nitrogen in a mortar together with 100 mg of dry Polyclar AT (insoluble polyvinylpyrrolidone). The resulting leaf powder was mixed with 5 ml of extraction medium contained in a 12 ml polypropylene centrifuge tube and heated in a water bath for 1 h to 65 °C, inverting the tube from time to time. The extraction medium consisted of 2 % (w/v) hexadecyltrimethylammoniumbromide, 1.4 M NaCl, 20 mM EDTA-Na₂ and 1 % (w/v) PVP-40 in 100 mM Tris, adjusted to pH 8.0 with HCl; shortly before use, 0.2 % (v/v) 2-mercaptoethanol was added to the me-





Figure 1. Upper part: Outline map of Germany including the borderlines of the state of Rhineland-Palatinate. Lower part: Outline map of Rhineland-Palatinate including the positions of the main rivers and the cities of Koblenz and Mainz. The locations of the pedunculate oak populations investigated are framed. The regularly budding stands are located in the forest districts Hagenbach, Simmern, Speyer and Trippstadt, the late budding stand is growing in the forest district Mayen. Saplings of the mother stands Simmern and Mayen are growing in the forest district Neupfalz (for further details see Table 1).

		Location	Location			
Forest region Forest district		Sub-district Stand		Altitude a.s.l. (m)	(years 2001)	
Palatinate Rhine valley	Hagenbach ^{a)}	Streitwald, south	VI 2 a ²	120	177	
Eifel	Mayen ^{b)}	Graf von Eltz ^{d)}	Abt. 28 C ¹	240-260	92	
Hunsrück	Simmern ^{c)}	Argenthal	13 a	430	196	
Palatinate Rhine valley	Speyer ^{c)}	Mutterstadt	1 7 '	100	123	
Palatinate Forest	Trippstadt ^{a)}	Langeck,	XI 15 b ²	410	170	
		Schmetzeck	XI 6 a ³			
Hunsrück	Neupflaz ^{a)}	Thiergarten	137 3 ^{e)}	420	12 ^{g)}	
Hunsrück	Neupflaz ^{a)}	Thiergarten	137, 7 ^{f)}	420	12 ^{g)}	

Table 1. Characterization of the seven pedunculate oak (*Quercus robur L.*) populations studied in the state of Rhineland-Palatinate, south west Germany.

^{a)} state forest, ^{b)} private forest, ^{c)} community forest, ^{d)} late budding provenance imported as seeds from Slovenia, ^{e)} offsprings of the stand investigated at forest district Simmern, ^{f)} offsprings of the stand investigated at forest district Mayen, ^{g)} the offsprings were 8 years old when investigated, they are part of a provenance trial established at forest district Neupfalz. In the sub-district Langeck 33 oak trees were probed while in the sub-district Schmetzeck short twigs with young leaves were sampled from 16 trees. At all other places 50 trees were probed.

dium. Afterwards, 1 ml of aqueous 5 M ammonium acetate solution and 6 ml of chloroform / isoamylalcohol (24 : 1 (v/v)) were added and the tube inverted several times. The mixture was centrifuged at 6 000 x g for 10 min (Sorvall RC-5B centrifuge) at RT. The upper aqueous phase, containing the DNA, was carefully taken off and transferred into another 12 ml centrifuge tube. Afterwards, 1 vol (v/v) of chloroform / isoamylalcohol (24:1 (v/v)) was added to further free the DNA from impurities, inverted several times and then centrifuged (as above). The aqueous phase was transferred to another 12 ml centrifuge tube, 2/3 vol(v/v) of cold (-20 °C) isopropanole added and the tube, carefully inverted, several times. Then it was stored for 30 min at -20 °C and centrifuged for 3 min at 10 000 x g and 4 °C. The supernatant was decanted, the pellet dried for 30 min in vacuum at RT and finally suspended in 1 ml of TE buffer (10 mM Tris, 1 mM EDTA-Na₂, adjusted to pH 8.0 with HCl). After addition of 3.3 μ l RNase A (3 $\mu g/\mu l$, Sigma) the suspension was incubated for 60 min at 37 °C and 1 ml of 5 M ammonium acetate in distilled water and 2.5 vol of pre-chilled (-20 °C) ethanol added. The DNA was precipitated after several careful inversions of the centrifuge tube and sedimented by centrifugation at 10 000 \times g for 10 min at 4 °C. The supernatant was decanted, the DNA precipitate washed with 3 ml of 80 % (v/v) aqueous ethanol, and centrifuged as before. The supernatant was decanted and the precipitate dried for 30 to 60 min in vacuum at RT. Finally the DNA was suspended in 500 µl TE-buffer and stored at 4 °C (or -20 °C) until used.

Estimation of the concentration and purity of DNA

The DNA concentration of leaf extracts was estimated photometrically according to the equation: c_{DNA} [µg/ml] = $(A_{260} - A_{320}) \times 50 \times \text{diluting factor (SAMBROCK et al.})$ 1989). DNA purity was determined from the ratio A_{260} $-A_{320}/A_{280} - A_{320}$ (SAMBROCK et al. 1989). The presence of high molecular weight DNA was determined by agarose electrophoresis and only extracts containing high molecular DNA were used for PCR reactions. Before electrophoresis DNA extracts were diluted to 1 μ g/5 μ l. To 5 μ l extract 2.5 μ l loading buffer (5 M urea and 0.02 % Xylene cyanole, BASSAM et al. 1991) was mixed and 5 µl of this mixture loaded per gel trough. Electrophoresis was performed for 90 min at RT and 100 V on 0.8 % agarose gels in TBE buffer (45 mM Tris, 40 mM boric acid, 1.25 mM EDTA-Na₂, pH 8.4). As size marker a Hind III restricted DNA of λ -phage (600 µg/ml, Eurogentec, Seraing, Belgium) was used. DNA extracts were stored at 4 °C, reagents at -20 °C.

PCR (polymerase chain reaction) procedure

The polymerase chain reaction (PCR) was performed in 15 ml reactions containing 7.5 μ l of mastermix and 7.5 μ l of DNA extract (10 ng/ μ l). Thermal cycling was done in a programmable heating block (MJ Research, type PTC-200, Watertown, Mass., USA). Reactions were started at 94 °C for 4 min, continued with 40 cycles at 94 °C for 45 sec, 36 °C for 45 sec (primer number 26, 28 and 29) or 46 °C for 45 sec (primer number 50 and 51) and 72 °C for 90 sec and terminated at 72 °C for 5 min.

The master mix consisted of $1.5 \,\mu l \,MgCl_2 \,(25 \,mM)$,

1.2 μ l sterile double distilled water, 1.5 μ l of dNTPs (2 mM each of dATP, dCTP, dGTP and dTTP), 1.5 μ l Stoffel-buffer (100 mM KCl, 100 mM Tris, adjusted with HCl to pH 8.3), 0.3 μ l DNA polymerase (10 U/ μ l, Stoffel fragment, Perkin Elmer/Cetus, Langen, Germany) and 1.5 μ l primer (30 μ M).

Primer sequences

Of several RAPD primers used to amplify oak DNA (MOREAU 1993, MOREAU *et al.* 1994) three were used in this study: primer number 26 (5'-ACGGATCCTG-3'), number 28 (5'-ACGGTACCAG-3') and number 29 (5'-TGCTGCAGGT-3') (WERNER *et al.* 1997). Of several anchored simple sequence repeat primers (ZIETKIEWICZ *et al.* 1994, WEISING *et al.* 1991) two brought interpretable DNA patterns: number 50 (5'-GTG(GATA)₄-3') and number 51 (5'-GGA(GATA)₄-3'). Primers were obtained from GENterprise, University of Mainz.

Agarose electrophoresis

Amplified DNA fragments were separated and visualized by electrophoresis in 1.2 % agarose gels in TBE buffer (45 mM Tris, 40 mM boric acid, 1.25 mM EDTA-Na₂), pH 8.4. A sample of 7 μ l amplified DNA was mixed with 3.5 μ l of loading buffer consisting of 5 M urea and 0.02 % Xylene cyanole and 8 μ l loaded per slot. As molecular marker a "Biomarker Low standards" comprising DNA fragments of 50, 100, 200, 300, 400, 500, 525, 700 and 1000 bp (600 μ g/ml, BioVentures, Eurogentec, Darmstadt, Germany) was used. The marker was mixed with an equal volume of loading buffer and 8 μ l applied per slot. Electrophoresis was performed for 4 h under a field strength of 100 V at RT. DNA bands were stained for 30 min in ethydiumbromide (5 μ g/ml) and destained for 30 min in tap water. Gels were photographed through a yellow filter under UV light (302 nm) using a Polaroid 667 film.

Evaluation of DNA-patterns

Photographs were used for analysis of the amplification products. The length of a DNA-amplification product was calculated from its position in the agarose gel relative to the standardised scale of DNA-marker bands. Amplification was repeated at least twice and the coarseness of each band determined. Only the bands reproducible in multiple runs, regardless of their intensity, were utilized in the survey (Table 2).

Quantification of genetic variation

Once RAPD- and AMP-loci were noticed in all investigated individuals the presence (1) or absence (0) of phenotypes was entered into a 1/0-matrix. A RAPD-/AMP-locus was considered to be polymorphic if its phenotypes varied.

The genetic difference between two populations was taken as equivalent to the difference in their RAPD or AMP patterns and quantified by the Tanimoto distance, δ_T , using the computer programm "CoTrix, a computer

Table 2. Fragment length of 64 loci, amplified with three RAPD primers (number 26, 28 and 29) and two AMP primers (number 50 and 51).

no. 26	no. 28	no. 29	no. 50	no. 51
A (120±10)	A (120±10)	A (200±10)	A (220±10)	A (220±10)
B (180±10)	B (180±10)	B (270±10)	B (260±10)	B (260±10)
C (230±10)	C (220±10)	C (350±10)	C (350±10)	C (300±10)
D (300±10)	D (280±10)	D (400±15)	D (400±10)	D (350±10)
E (370±10)	E (330±10)	E (470±15)	E (430±10)	E (400±10)
F (400±10)	F (390±10)	F (600±15)	F (470±10)	F (430±10)
G (430±10)	G (440±10)	G (680±15)	G (500±10)	G (470±10)
H (470±10)	H (520±15)	H (830±20)	H (580±15)	H (500±10)
I (520±15)	I (600±15)	I (900±20)	I (630±15)	I (580±10)
J (620±15)	J (660±20)	J (950±20)	J (700±15)	J (630±10)
K (650±15)	K (740±20)	K (1050±20)	K (750±15)	K (680±10)
L (730±20)	L (850±20)		L (930±20)	L (700±10)
M (860±20)			M (1000±20)	M (750±15)
				N (930±15)
				O (1000±20

programme for analysis of DNA banding patterns, version 2.0" (DEGEN 1996).

Relative frequencies of homozygous null genotypes (0/0) were calculated and then used to determine allele frequencies under the assumption that RAPD and AMP genotypes are two allelic systems and that genotypes are distributed according to Hardy-Weinbergs' theorem (STEPHENS *et al.* 1992, LYNCH AND MILLIGAN 1994, DAWSON *et al.* 1995). Accordingly, frequencies of the null allele (*p*) resulted from $p = \sqrt{0/0}$ and those of the one allel (*q*) from q = 1 - p. Allel frequencies were calculated by use of the computer program "Population Genetic Analysis" (PopGene, Version 1.21, YEH *et al.* 1997).

Expected heterozygosities (H_l) , (H_e) and (H_s) were calculated by use of the computer program PopGene, version 1.21 (YEH *et al.* 1997) as were Nei's total genetic diversity (H_T) , diversity within populations (H_s) , diversity between populations (D_{ST}) and the coefficient of the genetic diversity (G_{ST}) (NEI 1973, 1987).

Cluster analyses were performed following the unweighted pair-group method with arithmetic mean (UPGMA) (SNEATH & SOKAL 1973, FERGUSON 1980).

RESULTS

Amplification success

On average 88 % (between 50 to 96 %) of all investigated pedunculate oaks resulted in clearly identifiable DNA patterns. Amplification of random DNA was primer dependent and population specific. The least amplification success was obtained with trees of the Trippstadt population (RAPD: 68 %, AMP-PCR: 52 %), the best success was obtained with individuals of the offspring of the stands Mayen and Simmern (RAPD: 93 %, AMP-PCR: 93 %).

Number and size of amplified DNA fragments

By use of primers number 26, 28 and 29, at maximum 13, 12 and 11 DNA bands could be accessed. Their sizes ranged from 1050 to 120 base pairs (Table 2). When using the anchoring microsatellite primers No. 50 and 51, a maximum of 13 and 15 DNA bands could be accessed, respectively. Their fragment length ranged from 1000 to 220 bp (Table 2).

Polymorphic loci

Of the 64 loci which were accessible on average 87 % were polymorphic. Of the RAPD primer loci 64 to 100 % and of the AMP primer loci 69 to 100 % proved to be polymorphic. The number of polymorphic loci decreased in the two offspring populations as compared to their parent populations. The decrease was 6 % in offspring of the regularly budding population Simmern while it was 8 % in offspring of the late budding population Mayen.

Tanimoto distances

The Tanimoto-distance, which can be applied to dominant DNA markers like RAPD and AMP markers, was used to calculate the sub-population differentiation which equals the genetic difference between a certain sub-population and the complement of the remaining sub-populations.

The average sub-opulation Tanimoto-distance was exclusively related to the five adult populations, $\delta_T = 46.18$ %. The smallest distance was observed for the regularly budding stand Trippstadt ($\delta_T = 44.71$ %), the largest distance was calculated for the late budding stand Mayen ($\delta_T = 47.42$ %). (At a value of 100 % all individuals would be unequal.)

Table 3. Tanimoto-distances of the 7 pedunculate oak populations under study in relation to the remaining complement of the six populations (sub-population Tanimoto-distances).

Population	Average sub-population Tanimoto-distance at the gene cluster								
	no. 26	no. 28	no. 29	Mean	no. 50	no. 51	Mean	Grand mean	
Hagenbach	0.5792	0.4097	0.3802	0.4564	0.5786	0.3629	0.4708	0.4621	
Mayen	0.5071	0.4362	0.4928	0.4787	0.5187	0.4030	0.4609	0.4716	
Mayen-offspring	0.5213	0.4436	0.4257	0.4635	0.4408	0.3510	0.3959	0.4365	
Simmern	0.5040	0.4263	0.4385	0.4562	0.4612	0.3814	0.4213	0.4423	
Simmern-offspring	0.5490	0.4785	0.4375	0.4883	0.4205	0.3990	0.4098	0.4569	
Speyer	0.5420	0.3950	0.4836	0.4735	0.5492	0.4093	0.4793	0.4758	
Trippstadt	0.5557	0.4442	0.4082	0.4694	0.4343	0.3800	0.4072	0.4445	
Grand mean	0.5369	0.4334	0.4366	0.4694	0.4862	0.3839	0.4350	0.4554	

© ARBORA PUBLISHERS

O. WERNER & G. M. ROTHE: GENETIC DISTANCE AND TRANSMISSION AT RAPD AND SSR LOCI IN OAK



Figure 2a. DNA patterns of DNA regions amplified with the AMP-primer No. 50. Lanes 1 and 22: DNA size standard. Lanes 2 to 21: DNA of trees number 22 to 42 of stand Hagenbach.



Figure 2b. DNA patterns of DNA regions amplified with th AMP-primer No. 51. Lanes 1 and 22: DNA size standard Lanes 2 to 21: DNA of offsprings number 21 to 40 of the Simmern stand.

Table 4. Expected heterozygosities H_e at the loci amplified with three RAPD primers (number 26, 28 and 29) and two AMP primers (number 50 and 51).

Population –	Expected heterozygosities (H_e) of primer loci							
	no. 26	no. 28	no. 29	no. 50	no. 51	Mean (H_e)		
Hagenbach	0.2559	0.3601	0.2297	0.2952	0.1839	0.2650		
Mayen	0.2906	0.3705	0.3208	0.2612	0.2187	0.2924		
Mayen-offspring	0.3020	0.2538	0.2587	0.2517	0.1280	0.2388		
Simmern	0.3111	0.4123	0.2772	0.2447	0.2365	0.2964		
Simmern-offspring	0.3418	0.3412	0.2673	0.1488	0.2364	0.2671		
Speyer	0.3578	0.3171	0.3290	0.2922	0.2463	0.3085		
Trippstadt	0.2783	0.3016	0.2717	0.1728	0.1913	0.2430		
Mean (H_s)	0.3054	0.3367	0.2792	0.2381	0.2059	0.2730		
Mean		0.3071		0.2	220			

When distances were calculated including the two offspring populations then the average subpopulationdistance over all seven populations was $\delta_T = 45.57$ %. The smallest distance resulted now for the offspring of the late budding stand Mayen ($\delta_T = 43.65$ %) while the largest distance was observed for the regularly budding stand Speyer ($\delta_T = 47.58$ %) (Table 3, Figure 3).

In comparing the two parent stands Simmern and Mayen with each other and with their offspring populations deviating trends were observed. The subpopulation differentiation was larger in the offspring of the Simmern stand than in the Simmern parent stand by 1.46 %, however the distance decreased in offspring of the Mayen stand in comparison to the Mayen parent stand by 3.51 % (Table 3, Figure 3). The maximum difference between subpopulation distances was 3.93%. The genetic distances of gene loci amplified with RAPD primers were on average a little larger (δ_T = 46.94 %) than the ones amplified with AMP primers (δ_T = 43.50 %) (Table 3).

Expected heterozygosities

The average expected heterozygosity of the five adult stands was $H_e = 28.11$ %, the smallest and the largest value were found for the regularly budding stands Trippstadt ($H_e = 24.30$ %) and Speyer ($H_e = 30.85$ %), respectively (Table 4). Including the two offspring populations, the mean expected heterozygosity was $H_e = 27.30$ %, with the smallest expected heterozygosity resulting for the Mayen offspring population ($H_e = 23.88$ %) and the largest for the Speyer stand ($H_e = 23.88$ %)



Figure 3. Tanimoto sub-population distances related to the 64 loci investigated. : average of all seven distances.



Figure 4. Tanimoto sub-population distances related to the loci amplified with primers number 26, 28 and 29. δ : average of the seven distances.

30.85 %) (Table 4).

Heterozygosity was reduced from the parent stands to the offspring populations. From the regularly budding parent stand Simmern to their offspring heterozygosity decreased by 2.93 % and in offspring from the late budding Mayen stand heterozygosity decreased by 5.36 % (Table 4). With RAPD loci, the heterozygosity of the Simmern offspring population decreased in relation to the parent population by 1.67 % and in the Mayen population it decreased in relation to the parent population by 5.58 %. With AMP amplification, the heterozygosity of the Simmern parent and offspring

© ARBORA PUBLISHERS

Tanimoto subpopulation distance, primer 50-51



Figure 5. Tanimoto sub-population distances related to the loci amplified with primers number 50 and 51. : average of the seven distances.

populations decreased by 4.80 % and in the Mayen parent and offspring populations it decreased by 5.01 % (Table 4).

Genetic diversity

The genetic diversity H_T (NEI 1973) of the studied populations was primarily based upon the genetic diversity within the populations. The genetic diversity between populations (D_{ST}) in relationship to the total genetic diversity (H_T) was only 9.06 % (Table 5). As a result, the genetic diversity within populations was 90.94 %. The G_{ST} value related to the gene loci amplified with RAPD primers was a little smaller ($G_{ST} = 8.94$ %) than the G_{ST} value ($G_{ST} = 9.25$ %) of the samples analyzed with the AMP primers (Table 5).

DISCUSSION

RAPD markers have been increasingly applied to provide estimates of genetic diversity within and between tree populations (AIDE & RIVERA 1998, ALLNUTT *et al.* 1998, 1999, CHALMERS *et al.* 1992, GILLIES *et al.* 1997, NEWTON *et al.* 1999). Of the six decameric RAPD primers (MOREAU 1993, MOREAU *et al.* 1994) tried in this study to quantitate genetic variation in pedunculate oak, three (Nos. 26, 28 and 29) resulted in reproducible and interpretable DNA patterns.

SSR motives, such as $(CA/GT)_n$ and $(GA/CT)_n$ were investigated in five tropical tree species and found to be relatively abundant with 10⁴ and 10⁵ such sites in the genomes (CONDIT & HUBBELL 1991). In bur oak (*Q. macrocarpa*), the number of dinucleotide repeats per genome was estimated to be 3 x 10⁵ (BARRENECHE *et al.* 1998). The (GA/CT)_n repeats were sequenced and showed average repeat length of 16.2 (Dow *et al.* 1995). In *Quercus robur* and *Q. petraea* the genome

Diversity measure	Genetic diversity of primer loci								
	no. 26	no. 28	no. 29	no. 50	no. 51	Mean			
H_T H_S	0.3401 0.3053	0.3643 0.3367	0.3148 0.2792	0.2831 0.2419	0.2277 0.2059	0.3060 0.2738			
G _{ST}	0.1014	0.0699	0.0970	0.1079	0.0770	0.0906			
		0.0894			0.09245				

Table 5. Nei's genetic diversities.

Table 6. Relative band-frequencies (p%) at RAPD and AMP loci which differ in the two parent stands and their offsprings by more than 30%.

Locus	Mayen				Simm		
	parent stand p (%)	progeny p (%)	Δ (%)	Locus	parent stand p (%)	progeny p (%)	Δ(%)
26-K	78	46	-32	26-K 26-L	81 67	40 27	-41 -40
28-B	47	5	-42	28-B	67	16	-51
28-C	42	9	-33				
28-D	81	48	-33				
28-F	63	25	-38				
28-H	44	93	49				
29-I	41	72	31				
50-A	27	94	67				
50-E	18	60	42				
51-C	46	13	-33				
51-H	67	98	31				
				51-F	55	24	-1

Mayen: 7 RAPD and 4 AMP loci with band frequencies > 30 %. Simmern: 3 RAPD and 1 AMP locus with band frequencies > 30 %, Tanimoto sub population differences (δ_r): Mayen = 47.16 %, Mayen – offspring = 43.65 %, Simmern = 44.23 %, Simmern – offspring = 45.69 %. Pairwise Tanimoto distances (δ_r): Mayen / Simmern = 45.45 %, Mayen – offspring / Simmern – offspring = 40.31 %, Mayen – offspring = 46.21 %, Simmern / Simmern offspring = 42. 90 %. Simmern: regularly flushing pedunculate oak, Mayen: late flushing pedunculate oak.

size is 0.86×10^9 bp in the haploid genome with about one (GA)_n site every 120 kb (STEINKELLNER *et al.* 1997, BARRENECHE *et al.* 1998). Investigating genetic polymorphisms of 5 GA-microsatellite DNA loci 5 to 12 alleles per locus were observed in *Q. petraea* and *Q. robur* (STEINKELLNER *et al.* 1996). In both species, the same level of differentiation was detected and none was restricted to one species alone. In these studies microsatellite sequences were amplified by PCR using primers from the conserved flanking regions (STEINKEL-LNER *et al.* 1996).

Anchored microsatellite primed (AMP)-polymerase chain reaction (PCR), or as it is also named, simple sequence repeat anchored (SSRA)-PCR, amplifies genomic segments flanked by inversely oriented, closely spaced, repeated sequences (ZIETKIEWICZ *et al.* 1994). AMP-PCR depicts various repetitive sequences in eucariotic genomes with a length of two to six bases (ZIETKIEWICZ et al. 1994, WINTER & KAHL 1995). As primers, oligonucleotides are used that are composed of a complementary repetitive sequence and a random two to four nucleotide long sequence at the 5' terminus. This end anchors outside the repetitive sequence within the genome (ZIETKIEWICZ et al. 1994). Polymorphisms evolve at RAPD and SSR loci either by mutation (insertion, deletion), within sequences between primer binding sites, or by alterations within the primer binding sites themselves (WINTER & KAHL 1995). AMP-PCR is similar to RAPD-PCR in that it also uses only one primer, needs no sequence information prior to analysis, and its amplified sequences are dominantly inherited (TROGGIO et al. 1996, ZIETKIEWICZ et al.

Locus	Mayen				Simn		
	parent stand H_e (%)	offspring H_e (%)	Δ (%)	Locus	parent stand H_e (%)	offspring H_e (%)	Δ (%)
26-E	0	49.92	49.92				
28-B	39.29	4.49	-34.80	28-B	48.80	14.90	-33.90
28-K	49.06	0	-49.06				
29-D	46.49	0	-46.49	29-D	47.46	0	-47.46
				29-F	0	32.75	32.75
				50-G	49.64	0	-49.64
51-A	45.97	0	-45.97				
51-D	35.03	0	-35.03				
				51-E	34.12	0	-34.12

Table 7. Expected heterozygosities (H_{ϵ} %) at RAPD and AMP loci which differ in the two studied pedunculate oak stands and their offsprings by more than 30 %.

Mayen: 4 RAPD and 2 AMP loci with expected heterozygosities > 30 %. Simmern: 3 RAPD and 2 AMP loci with expected heterozygosities > 30 %. Simmern: Regularly flushing oak. Mayen: Late flushing oak.

1994). Here we tried 24 anchored microsatellite primers

of the types

5'-NNN(GTG)₅-3' and 5'-NNN(GATA)₄-3', which resulted in sharp and reproducible bands. Among these, the latter type revealed the greater differentiation.

The average number of polymorphic RAPD loci studied in this investigation was 89.57 % (64 to 100 %). In various tree species this number ranges from 35.7 %(Pigerodendron uviferum (ALLNUTT et al. 1999)) to 100 % (Picea mariana (ISABEL et al. 1995)) and on average (8 species) it is 71.34 % (ALLNUTT et al. 1999). The average number of polymorphic AMP loci which we investigated was 85.07 % (69 to 100 %). Thus the average number of polymorphic AMP loci was a little lower (-3.5 %) than the number of RAPD loci. The chance that RAPD primers amplify SSR loci is rather low and may be one in twenty, as estimated for Scots pine (LU et al. 1997). In RAPD, the absence of an amplification product can mostly be traced back to an alteration of the primer binding site, as shown in Pinus sylvestris (LU et al. 1997).

During the re-immigration of sessile oak and pedunculate oak from the Mediterranean area to Central Europe after the last ice age, several subpopulations evolved. They differ in a certain sequence of the plastidic t-RNA-gene (KREMER *et al.* 1991; PETIT *et al.* 1993). However, small area population differences cannot be elucidated with these sequences.

Although the genetic distances between the pedunculate oak populations which we investigated here are low ($G_{ST} \le 10$ %) they can be differentiated according to frequency differences at RAPD and AMP primed loci. Frequencies of Tanimoto's sub-population dis-

tances depended on the DNA sequences investigated. Within RAPD loci amplified with primers No. 26, 28 and 29 sub-population distances increased as follows: Simmern (45.46 %), Hagenbach (45.53 %), Trippstadt (45.53 %), Speyer (46.62) and Mayen (47.25 %), with the late budding population Mayen having the largest distance. With the AMP loci amplified with primers No. 50 and 51 sub-population distances increased differently within the investigated stands: Trippstadt (43.59 %), Simmern (44.48 %), Hagenbach (47.59 %), Mayen (47.68 %) and Speyer (48.53 %). Here, the late budding stand came fourth before the reglarly budding population at Speyer. At all 64 loci together subpopulation differences came close to those obtained with the two AMP loci: Trippstadt (44.71 %), Simmern (45.07 %), Hagenbach (46.31 %), Speyer (47.38 %) and Mayen (47.42 %). Here again the late budding variety exhibited the largest distance towards the remaining complement of the four populations.

The investigated parent populations Mayen and Simmern are certified for seed harvesting according to the official directives for forest reproductive materials in Germany ("Gesetz über forstliches Saat- und Pflanzgut"). The offspring of both stands were raised in a nursery for three years and then transplanted to a different forest district where they are part of a provenance trial. Sub-population distances in both offspring are different from those of the respective parental stands. In offspring of the Simmern stand, sub-population differentiation was 1.46 % larger than in the parent stand while in offspring of the late budding stand Mayen it was 3.71 % lower than in the parent stand. Moreover, at several single loci there are clear differences in both offspring (Table 6). In offspring of the late budding Mayen stand at 11 different loci frequency differences of 31 % and more were observed in comparison to the parent trees. At six loci (26-K, 28-B, -C, -D, -F and 51-C), the frequencies were lower in the offspring than in the parent trees, while at five loci (28-H, 29-I, 50-A, 50-E and 51-H) they were larger (Table 6). These alterations may be considered as a selection towards the climatic and edaphic conditions at the Hunsrück hills as compared to the Slovenian habitat. In offspring of the regular budding stand Simmern, at four different loci (26-K, 26-L, 28-B and 51-F), frequencies decreased by more than 30 % as compared to the parent trees (Table 6). These alterations occurred predominantly at RAPD-loci (Maven: 7 RAPD and 4 AMP loci. Simmern 3 RAPD and 1 AMP locus). That RAPD-loci may be subjected to selection has been demonstrated for Limber pine (Pinus flexilis James) (LATTA & MITTON 1997).

Different genetic developments in the two offspring populations investigated can also be recognized when expected heterozygosities are compared. The average expected heterozygosity calculated for stand Mayen was He = 29.24 % and that of its offspring He = 23.88 %. For the Simmern stand, the expected heterozygosity was He = 29.64 % and that of its offspring He = 26.71 %. BARRENECHE et al. (1998) found 18 % of RAPD markers to be heterozygous in two Q. robur trees. The reason for the reduction in the expected heterozygosities in offspring of the Mayen stand (-5.36 %) and in the Simmern stand (-2.93 %) may be that seeds were collected from a inadequate number of mother trees. A loss in genetic variation is generally thought to infer adaptive abilities of a population, but heterozygosity may increase during stand growth in case of selection against homozygotes.

Single locus events can also be recognized when expected heterozygosities are compared between the offspring and parental populations (Table 7). In the offspring of the late budding Mayen stand, expected heterozygosity decreased at 5 loci by more than 35 % (28-B, -K, 29-D, 51-A, 51-D) and increased by more than 35 % at one locus (26-E) as compared to the parental stand (Table 7). In the offspring of the regularly budding Simmern stand, the expected heterozygosity decreased at four loci by at least 33 % (28-B, 29-D, 50-G and 51-E), but increased at one locus by 33 % (29-F) as compared to the parental stand.

Altogether, clear variations in the two offspring generations are indicated. In the late budding Mayen population, frequency of alterations at 10 out of 64 loci (= 16 %) can be taken as significant, while in the regular budding Simmern population, serious frequency alterations affected only 4 out of 64 loci (= 6%). These, and lower frequency differences contribute to

the unequal population differentiations. To avoid a further reduction in genetic variability, stands should be created with young plants from different sources but similar habitat.

ACKNOWLEDGEMENT

Financial support of the "Forschungsanstalt für Waldökologie und Forstwirtschaft Rheinland-Pfalz" and the "Zentrum für Umweltforschung" of Mainz University is acknowledged and greatly appreciated.

LITERATUR

- AAS, G. 1991: Kreuzungsversuche mit Stiel- und Traubeneichen (Quercus robur L. und Quercus petraea [Matt.] Liebl.). Allgem. Forst- und Jagdzeitung 162: 462–465.
- AIDE, T. M. & RIVERA, E. 1998: Geographic patterns of genetic diversity in *Poulsenia armata (Moraceae)*: implications for the theory of Pleistocene refugia and the importance of riparian forest. J. Biogeography 25: 695-705.
- ALLNUTT, T. R., THOMAS, P., NEWTON, A. C. & GARDNER, M. F. 1998: Genetic variation in *Fitzroya cupressoides* cultivated in the British Isles, assessed using RAPDs. *Bot. J. Edinburgh* 55 (3): 329–341.
- ALLNUTT, T. R., NEWTON, A. C., LARA, A., PREMOLI, A., ARMESTO, J. J., VERGARA, R. & GARDNER, M. 1999: Genetic variation in *Fitzroya cupressoides (alerce)*, a threatened South American Conifer. *Molecular Ecology* 8: 975–987.
- BARRENECHE, T., BODENES, C., LEXER, C., TRONTIN, J.-F., FLUCH, S., STREIFF, R., PLOMION, C., ROUSSEL, G, STEINKELLNER, H., BURG, K., FAVRE, J.-M., GLÖSSL, J. & KREMER, A. 1998: A genetic linkage map of *Quercus* robur L. (pedunculate oak) based on RAPD, SCAR, microsatellite, minisatellite, isozyme and 5 S rDNA markers. *Theor. Appl. Genet.* 97: 1090–1103.
- BASSAM, B. J., CAETANO-ANOLLÉS, G. & GRESSHOFF, P. M. 1991: Fast and sensitive silver staining of DNA in polyacrylamide gels. *Analyt. Biochem.* 196, 80–83.
- CHALMERS, K. J., WAUGH, R., SPRENT, J. I., SIMONS, A. J. & POWELL, W. 1992: Detection of genetic variation between and within populations of *Gliricidia sepium* and *G maculata* using RAPD markers. *Heredity* 69: 465–472.
- CHALMERS, K., NEWTON, A. C., WAUGH, R., WILSON, J. & POWELL, K. 1994: Genetic variation in mahoganies (*Meliaceae*) detected using RAPDs. *Theoretical and Applied Genetics*: 89 (4): 504–508.
- CONDIT, R. & HUBBELL, S. P. 1991: Abundance and DNA sequence of two-base repeat regions in tropical tree genomes. *Genome* 34: 66–71.
- DAWSON, J. K., SIMONS, A. J., WAUGH, R. & POWELL, W. 1995: Diversity and genetic differentiation among subpopulations of *Gliricidia sepium* revealed by PCRbased assays. *Heredity* 74: 10–18.
- DEGEN, B. 1996: CoTrix, a computer programme for analysis of DNA banding patterns – Version 2.0. Institut für Forstgenetik, Grosshansdorf (http://www.rrz.unihamburg.de/OekoGenetik/welcome.htm).

- DELLAPORTA, S. L., WOOD, J. & HICKS, J. B. 1983: Plant DNA minipreparation: version II. *Plant Molecular Biology Reporter* 1: 19–21.
- DOW, B. D., ASHLEY, M. V. & HOWE, H. F. 1995: Characterization of highly variable (GA/CT)_n microsatellites in the bur oak, *Quercus macrocarpa. Theor. Appl. Genet.* 91: 137–141.
- DOYLE, J. 1991: DNA protocols for plants. Molecular techniques in taxonomy. Springer-Verlag, 283–293.
- FERGUSON, A. 1980: Biochemical systematics and evolution. Blackie, Glasgow and London.
- GILLIES, A. C. M., CORNELIUS, J. P., NEWTON, A. C., NAVAR-RO, C., HERÁNDEZ, M. & WILSON, J. 1997: Genetic variation in Costa Rican populations of the tropical timber tree *Cedrela odorata* L. (Spanish cedar), assessed using RAPDs. *Molecular Ecology* 6: 1133–1145.
- GILLIES, A. C. M., NAVARRO, C., LOWE, A. J., NEWTON, A. C., HERNÁNDEZ, M., WILSON, J. & CORNELIUS, J. P. 1999: Genetic diversity in Mesoamerican populations of mahogany (*Swietenia macrophylla*), assessed using RAPDs. *Heredity* 83: 722–732.
- ISABEL, N., BEAULIEU, J. & BOUSQUET, J. 1995: Complete congruence between genetic diversity estimates derived from genotypic data at enzyme and random amplified polymorphic DNA loci in black spruce. *Proc. Nat. Acad. Sci., USA*, 92: 6369–6373.
- KREMER, A., PETIT, R. J., ZANETTO, A., FOUGERE, V., DU-COUSSO, A., WAGNER, D. & CHAUVIN, C. 1991: Nuclear and organelle gene diversity in *Quercus robur* and *Quercus petraea*. *In*: Müller-Starck, G. & Ziehe, M.: Genetic variation in european populations of forest trees. J. D. Sauerländer's Verlag, Frankfurt, 141–166.
- LATTA, R. G & MITTON, J. B. 1997: A comparison of population differentiation across four classes of gene marker in Limber pine (*Pinus flexilis James*). *Genetics* **146**: 1153 -1163.
- LU, M.-Z., WANG, X.-R. & SZMIDT, A. E. 1997: Molecular properties of RAPDs in *Pinus sylvestris* (L.) and their implications for genetic analysis. *For. Genet.* **4** (4): 227–234.
- LYNCH, M. & MILLIGAN, B. G. 1994: Analysis of population genetic structure with RAPD markers. *Mol. Ecol.* 3: 91–99.
- MOREAU, F. 1993: Recherche de marquers appliquée a la differenciation moleculaire entre chêne sessile (*Quercus petraea*) et pedonculé (*Quercus robur*). Thèse no. 593, Université de Poitiers, Faculté des Science Fondamentales et Appliquées.
- MOREAU, F., KLEINSCHMITT, J. R. G. & KREMER, A. 1994: Molecular differentiation between *Q. petraea* and *Q. robur* assessed by random amplified DNA fragments. *For. Genet.* 1(1): 51–64.
- NEI, M. 1973: Analysis of gene diversity in subdivided populations. Proc. Nat. Acad. Sci. USA 70: 3321–3323.
- NEI, M. 1987: Molecular Evolutionary Genetics. Columbia University Press, New York.
- NEWTON, A. C., ALLNUTT, T., GILLIES, A. C. M., LOWE, A. & ENNOS, R. A. 1999: Molecular phylogeography, intraspecific variation and the conservation of tree species. *Trends in Ecology and Evolution* **14** (4): 140–145.
- PETIT, R. J., KREMER, A. & WAGNER, D. B. 1993: Geographic structure of chloroplast DNA polymorphisms in European oaks. *Theor. Appl. Genet.* 87: 122–128.

- RÖHRIG, E. & GUSSONE, H. A. 1980: Waldbau auf ökologischer Grundlage: Der Wald als Vegetationstyp und seine Bedeutung für den Menschen. Verlag Paul Parey, Hamburg und Berlin.
- ROMMEL, M., ROTHE, G. M., MAURER, W. D. & TABEL, U. 1995: Artbestimmung bei Stiel- und Traubeneichen. *In*:
 W. D. Maurer & U. Tabel (Hrsg.) Genetik und Waldbau unter besonderer Berücksichtigung der heimischen Eichenarten. Mitteilungen aus der Forstlichen Versuchsanstalt Rheinland-Pfalz, Nr. 34/95, Trippstadt, S. 310 -320.
- SAMBROOK, J., FRITSCH, E. F. & MANIATIS, T. 1989: Molecular cloning, a laboratory manual. Cold Spring Harbor Laboratory Press. New York.
- SNEATH, P. H. A. & SOKAL, R. R. 1973: Numerical Taxonomy: The Principle and Practice of Numerical Classification. Freeman, San Francisco.
- STEINKELLNER, H., FLUCH, S., TURETSCHEK, E., STREIFF, R., KREMER, A., BURG, K. & GLÖSSEL, J. 1996: Development of microsatellite loci from *Quercus robur* and *Quercus petraea. In*: M. R. Ahuja *et al.* (eds.), Somatic Cell Genetics and Molecular Genetics of Trees, 217–222. Kluwer Academic Publishers, Netherlands.
- STEINKELLNER, H., FLUCH, S., TURETSCHEK, E., LEXER, C., STREIFF, R., KREMER, A., BURG, K. & GLÖSSEL, J. 1997: Identification and characterization of $(GA/CT)_n$ microsatellite loci from *Quercus petraea*. *Plant Mol. Biol.* **33**: 1093–1096.
- STEPHENS, J. C., GILBERT, D. A., YUHKI, N. & O'BRIEN, S. J. 1992: Estimation of heterozygosity for single-probe mulilocus DNA fingerprints. *Mol. Biol. Evol.* 9(4): 729–743.
- TROGGIO, M., DIMASSO, E., LEONARDI, S., CERONI, M., BUCCI, G., PIOVANI, P. & MENOZZI, P. 1996: Inheritance of RAPD and I-SSR markers and population parameters estimation in European beech (*Fagus sylvatica L.*). For. Genet. 3(4): 173–181.
- WEISING, K., RAMSER, J., KAEMMER, D., KAHL, G & EPPLEN, J. T. 1991: Oligonucleotide fingerprinting in plants and fungi. *In*: Burke, T., Dolf, G, Jeffreys, A. J. & Wolff, R. (eds.): DNA fingerprinting: approaches and applications. Birkhäuser Verlag, Basel, Boston, Berlin, 312–328.
- WERNER, O., ROTHE, G M., MAURER, W. D. & TABEL, U. 1997: Differentiation of pedunculate oak (*Quercus robur* L.) populations by DNA amplification fingerprinting. *In*: Steiner, K. C. (ed.): Diversity and adaptation in oak species, Proceedings, The Pennsylvania State University, Pennsylvania, USA. October 12–17. pp. 22–36.
- WINTER, P. & KAHL, G 1995: Molecular marker technologies for plant improvement. World Journal of Microbiology & Biotechnology, 11, 438–448.
- YEH, F. C., CHONG, D. K. X. & YANG, R.-C. 1995: RAPD variation within and among natural populations of trembling aspen (*Populus tremuloides* Michx.) from Alberta. *Journal of Heredity* 86: 454–460.
- YEH, F. C., YANG, R. C. & BOYLE, T. 1997: PopGene Version 1.21. Microsoft Window-based freeware for population genetic analysis (http://www.ualberta.ca/~fyeh).
- ZIETKIEWICZ, E., RAFALSKI, A. & LABUDA, D. 1994: Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20: 176–183.

© ARBORA PUBLISHERS